

For use in exams from the June 2016 Series onwards

## DATA - FUNDAMENTAL CONSTANTS AND VALUES

Quantity	Symbol	Value	Units
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$
magnitude of the charge of electron	$e$	$1.60 \times 10^{-19}$	C
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	$\alpha$	$2.90 \times 10^{-3}$	m K
electron rest mass (equivalent to $5.5 \times 10^{-4}$ u)	$m_e$	$9.11 \times 10^{-31}$	kg
magnitude of electron charge/mass ratio	$\frac{e}{m_e}$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$
proton rest mass (equivalent to 1.00728 u)	$m_p$	$1.67(3) \times 10^{-27}$	kg
proton charge/mass ratio	$\frac{e}{m_p}$	$9.58 \times 10^7$	$\text{C kg}^{-1}$
neutron rest mass (equivalent to 1.00867 u)	$m_n$	$1.67(5) \times 10^{-27}$	kg
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$
atomic mass unit (1u is equivalent to 931.5 MeV)	u	$1.661 \times 10^{-27}$	kg

### ALGEBRAIC EQUATION

quadratic equation  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

### ASTRONOMICAL DATA

Body	Mass/kg	Mean radius/m
Sun	$1.99 \times 10^{30}$	$6.96 \times 10^8$
Earth	$5.97 \times 10^{24}$	$6.37 \times 10^6$

### GEOMETRICAL EQUATIONS

arc length =  $r\theta$

circumference of circle =  $2\pi r$

area of circle =  $\pi r^2$

curved surface area of cylinder =  $2\pi r h$

surface area of sphere =  $4\pi r^2$

volume of sphere =  $\frac{4}{3}\pi r^3$

## Particle Physics

Class	Name	Symbol	Rest energy/MeV
photon	photon	$\gamma$	0
lepton	neutrino	$\nu_e$	0
		$\nu_\mu$	0
	electron	$e^\pm$	0.510999
	muon	$\mu^\pm$	105.659
mesons	$\pi$ meson	$\pi^\pm$	139.576
		$\pi^0$	134.972
	K meson	$K^\pm$	493.821
		$K^0$	497.762
baryons	proton	p	938.257
	neutron	n	939.551

## Properties of quarks

antiquarks have opposite signs

Type	Charge	Baryon number	Strangeness
<b>u</b>	$+\frac{2}{3}e$	$+\frac{1}{3}$	0
<b>d</b>	$-\frac{1}{3}e$	$+\frac{1}{3}$	0
<b>s</b>	$-\frac{1}{3}e$	$+\frac{1}{3}$	-1

## Properties of Leptons

		Lepton number
Particles:	$e^-, \nu_e; \mu^-, \nu_\mu$	+1
Antiparticles:	$e^+, \bar{\nu}_e, \mu^+, \bar{\nu}_\mu$	-1

## Photons and energy levels

*photon energy*  $E = hf = \frac{hc}{\lambda}$   
*photoelectricity*  $hf = \phi + E_{k(\max)}$   
*energy levels*  $hf = E_1 - E_2$   
*de Broglie Wavelength*  $\lambda = \frac{h}{p} = \frac{h}{mv}$

## Waves

*wave speed*  $c = f\lambda$      *period*  $f = \frac{1}{T}$   
*first harmonic*  $f = \frac{1}{2l} \sqrt{\frac{T}{\mu}}$   
*fringe spacing*  $w = \frac{\lambda D}{s}$      *diffraction grating*  $d \sin \theta = n\lambda$   
*refractive index of a substance s*,  $n = \frac{c}{c_s}$   
*for two different substances of refractive indices  $n_1$  and  $n_2$ ,*  
*law of refraction*  $n_1 \sin \theta_1 = n_2 \sin \theta_2$   
*critical angle*  $\sin \theta_c = \frac{n_2}{n_1}$  for  $n_1 > n_2$

## Mechanics

*moments*     *moment* =  $Fd$   
*velocity and acceleration*  $v = \frac{\Delta s}{\Delta t}$       $a = \frac{\Delta v}{\Delta t}$   
*equations of motion*  $v = u + at$       $s = \left(\frac{u+v}{2}\right)t$   
 $v^2 = u^2 + 2as$       $s = ut + \frac{at^2}{2}$   
*force*  $F = ma$   
*force*  $F = \frac{\Delta(mv)}{\Delta t}$   
*impulse*  $F \Delta t = \Delta(mv)$   
*work, energy and power*  $W = F s \cos \theta$   
 $E_k = \frac{1}{2} m v^2$       $\Delta E_p = mg\Delta h$   
 $P = \frac{\Delta W}{\Delta t}$ ,  $P = Fv$   
*efficiency* =  $\frac{\text{useful output power}}{\text{input power}}$

## Materials

*density*  $\rho = \frac{m}{V}$      *Hooke's law*  $F = k \Delta L$   
*Young modulus* =  $\frac{\text{tensile stress}}{\text{tensile strain}}$      *tensile stress* =  $\frac{F}{A}$   
*tensile strain* =  $\frac{\Delta L}{L}$   
*energy stored*  $E = \frac{1}{2} F \Delta L$

---

**Electricity**

*current and pd*       $I = \frac{\Delta Q}{\Delta t}$      $V = \frac{W}{Q}$      $R = \frac{V}{I}$

*resistivity*       $\rho = \frac{RA}{L}$

*resistors in series*       $R_T = R_1 + R_2 + R_3 + \dots$

*resistors in parallel*       $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$

*power*       $P = VI = I^2R = \frac{V^2}{R}$

*emf*       $\varepsilon = \frac{E}{Q}$        $\varepsilon = I(R + r)$

