



Rewarding Learning

**General Certificate of Secondary Education
2025**

Biology

Unit 2

Foundation Tier

[GBL21]

MONDAY 9 JUNE, MORNING

**MARK
SCHEME**

General Marking Instructions

Introduction

Mark schemes are intended to ensure that the GCSE examinations are marked consistently and fairly. The mark schemes provide markers with an indication of the nature and range of candidates' responses likely to be worthy of credit. They also set out the criteria which they should apply in allocating marks to candidates' responses.

Assessment objectives

Below are the assessment objectives for GCSE Biology.

Candidates must:

- AO1** demonstrate knowledge and understanding of: scientific ideas; and scientific techniques and procedures;
- AO2** apply knowledge and understanding of and develop skills in: scientific ideas; scientific enquiry, techniques and procedures; and
- AO3** analyse scientific information and ideas to: interpret and evaluate; make judgements and draw conclusions and develop and improve experimental procedures.

Quality of candidates' responses

In marking the examination papers, examiners should be looking for a quality of response reflecting the level of maturity which may reasonably be expected of a 16-year-old which is the age at which the majority of candidates sit their GCSE examinations.

Flexibility in marking

Mark schemes are not intended to be totally prescriptive. No mark scheme can cover all the responses which candidates may produce. In the event of unanticipated answers, examiners are expected to use their professional judgement to assess the validity of answers. If an answer is particularly problematic, then examiners should seek the guidance of the Supervising Examiner.

Positive marking

Examiners are encouraged to be positive in their marking, giving appropriate credit for what candidates know, understand and can do rather than penalising candidates for errors or omissions. Examiners should make use of the whole of the available mark range for any particular question and be prepared to award full marks for a response which is as good as might reasonably be expected of a 16-year-old GCSE candidate.

Awarding zero marks

Marks should only be awarded for valid responses and no marks should be awarded for an answer which is completely incorrect or inappropriate.

Marking calculations

In marking answers involving calculations, examiners should apply the 'own figure rule' so that candidates are not penalised more than once for a computational error.

Types of mark schemes

Mark schemes for tasks or questions which require candidates to respond in extended written form are marked on the basis of levels of response which take account of the quality of written communication.

Other questions which require only short answers are marked on a point for point basis with marks awarded for each valid piece of information provided.

Levels of response

Tasks and questions requiring candidates to respond in extended writing are marked in terms of levels of response. In deciding which level of response to award, examiners should look for the 'best fit' bearing in mind that weakness in one area may be compensated for by strength in another. In deciding which mark within a particular level to award to any response, examiners are expected to use their professional judgement. The following guidance is provided to assist examiners.

Threshold performance: Response which just merits inclusion in the level and should be awarded a mark at or near the bottom of the range.

Intermediate performance: Response which clearly merits inclusion in the level and should be awarded a mark at or near the middle of the range.

High performance: Response which fully satisfies the level description and should be awarded a mark at or near the top of the range.

Quality of written communication

Quality of written communication is taken into account in assessing candidates' responses to all tasks and questions that require them to respond in extended written form. These tasks and questions are marked on the basis of levels of response. The description for each level of response includes reference to the quality of written communication.

For conciseness, quality of written communication is distinguished within bands of response as follows:

Band A: Quality of written communication is excellent.

Band B: Quality of written communication is good.

Band C: Quality of written communication is basic.

In interpreting these level descriptions, examiners should refer to the more detailed guidance provided below:

Band A (Excellent): The candidate successfully selects and uses the most appropriate form and style of writing. Relevant material is organised with a high degree of clarity and coherence. There is widespread and accurate use of appropriate specialist vocabulary. Presentation, spelling, punctuation and grammar are of a sufficiently high standard to make meaning clear.

Band B (Good): The candidate makes a reasonable selection and use of an appropriate form and style of writing. Relevant material is organised with some clarity and coherence. There is some use of appropriate specialist vocabulary. Presentation, spelling, punctuation and grammar are sufficiently competent to make meaning clear.

Band C (Basic): The candidate makes only a limited selection and use of an appropriate form and style of writing. The organisation of material may lack clarity and coherence. There is little use of specialist vocabulary. Presentation, spelling, punctuation and grammar may be such that intended meaning is not clear.

When one response is required to gain a mark, candidates will not gain credit if a correct response is given alongside one or more incorrect responses. This is referred to as listing.

| | | | AVAILABLE MARKS |
|---|--|---|-----------------|
| 1 | <ul style="list-style-type: none"> adds fluid to sperm; testis; places sperm into vagina; scrotum; carries sperm to the urethra; | [5] | 5 |
| 2 | <p>(a) (i) 25 – 34;</p> <p>(ii) 75+; percentage increased between 2011–2020; from 7–9%/by 2%;</p> <p>(b) addictive/affects heart rate; reduces oxygen-carrying capacity of the blood; can cause lung cancer/bronchitis/emphysema;</p> | <p>[1]</p> <p>[3]</p> <p>[3]</p> | 7 |
| 3 | <p>(a) (i) histogram;</p> <p>(ii) 1.65 – 1.69;</p> <p>(iii) 2 + 6 + 7 + 5; 20;</p> <p>(b) mass; length of foot;</p> | <p>[1]</p> <p>[1]</p> <p>[2]</p> <p>[2]</p> | 6 |
| 4 | <p>(a) A – evaporation; B – diffusion;</p> <p>(b) cuticle; waterproof/waxy;</p> <p>(c) (i) close stomata;</p> <p>(ii) Any two from: support; transport; photosynthesis;</p> | <p>[2]</p> <p>[2]</p> <p>[1]</p> <p>[2]</p> | 7 |

| | | | AVAILABLE MARKS | |
|---|-------------------------|---|-----------------|----|
| 5 | (a) (i) | female condom; | [1] | 7 |
| | (ii) | 2.0 (correct data from table); $(2.0) \div 100 \times 100\,000 = 2000$; | [2] | |
| | (b) (i) | changes hormone levels; stops development of the ovum; | [2] | |
| | (ii) | sperm cannot reach eggs/no fertilisation; | [1] | |
| | (iii) | difficult to reverse; | [1] | |
| 6 | (a) | phagocytes; produce antibodies; | [2] | 10 |
| | (b) Any 2 pairs: | contain haemoglobin; carry/bind to oxygen; no nucleus; more space for haemoglobin/oxygen; biconcave shape; increase surface area; | [4] | |
| | (c) (i) | fibrinogen; | [1] | |
| | (ii) | traps blood cells; | [1] | |
| | (iii) | prevents blood loss; prevents infection; | [2] | |
| 7 | (a) | different form of the same gene; | [1] | 6 |
| | (b) (i) | a and a; A; Aa Aa aa aa; | [3] | |
| | (ii) | 50%; | [1] | |
| | (iii) | AA; | [1] | |

| | | | AVAILABLE MARKS | |
|----------|------------|---|--------------------|----|
| 8 | (a) | (i) tumour B encapsulated/has a capsule; tumour cells have not entered the bloodstream/not spreading; | [2] | 10 |
| | | (ii) malignant; | [1] | |
| | (b) | (i) 32 (correct data from table); correct calculation (e.g. 0.32×3150); 1008; | [3] | |
| | | (ii) total percentage of patients adds up to 126/greater than 100; | [1] | |
| | (c) | early detection; early treatment; increase chances of cure/increase survival rate; | [3] | |
| | | | | |

- 9 (a) (i) growth;
repair; [2]
- (ii) gametes; [1]
- (iii) ovaries;
testes; [2]
- (b) (i) 4; [1]
- (ii) (stage) A;
(stage) D; [2]

(iii) **Indicative content**

Changes that occurred between stage A and stage B:

1. nucleus/nuclear membrane disappears;
2. chromosomes duplicate (described);

Differences between mitosis and meiosis:

3. 2 daughter cells vs 4 daughter cells produced;
4. (mitosis produces) diploid cells (meiosis produces) haploid cells;
5. (mitosis produces) identical cells (compared to parent cell/each other) vs (meiosis produces) variation/genetically different cells;
6. (mitosis has) one cell division/no stage C vs (meiosis has) 2 divisions/stage C;

(comparison required in mp 3–6)

| Band | Response | Mark |
|------|--|---------|
| A | Candidates must use appropriate, specialist terms throughout to describe and explain their conclusions using at least 5 of the points . They use good spelling, punctuation and grammar and the form and style are of a high standard. | [5]–[6] |
| B | Candidates use some appropriate, specialist terms throughout to describe and explain their conclusions using at least 3 of the points . They use satisfactory spelling, punctuation and grammar and the form and style are of a satisfactory standard. | [3]–[4] |
| C | Candidates make little use of specialist terms throughout to describe and explain their conclusions using at least 1 of the points . The spelling, punctuation and grammar, form and style are of a limited standard. | [1]–[2] |
| D | Response not worthy of credit. | [0] |

[6]

AVAILABLE
MARKS

14

| | | | AVAILABLE MARKS |
|---------|---|--------------|-----------------|
| 10 (a) | A – aorta; B – pulmonary vein; F – vena cava; | [3] | |
| (b) | valve; prevents backflow of blood; into (left) atrium; | [3] | |
| (c) (i) | higher pressure; (comparison needed) | [1] | |
| (ii) | more oxygen in D; (comparison needed) less carbon dioxide in D; (comparison needed) (accept converse) | [2] | 9 |
| 11 (a) | bacteria; | [1] | |
| (b) | sexual intercourse ; | [1] | |
| (c) (i) | 1787 – 1684/103; (103 ÷ 1684) × 100; 6.116; 6.1; | [4] | |
| (ii) | increase in use of condoms ; fewer sexual partners; | [2] | |
| (d) | antibiotics; | [1] | 9 |
| | | Total | 90 |