

Surname	Centre Number	Candidate Number
First name(s)		0



GCSE

3430UE0-1



FRIDAY, 27 MAY 2022 – MORNING

SCIENCE (Double Award)

Unit 5 – CHEMISTRY 2

HIGHER TIER

1 hour 15 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	7	
2.	8	
3.	9	
4.	15	
5.	8	
6.	6	
7.	7	
Total	60	

3430UE01
01

ADDITIONAL MATERIALS

In addition to this examination paper you will need a calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink or ball-point pen. Do not use gel pen or correction fluid. You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

Question 7(a) is a quality of extended response (QER) question where your writing skills will be assessed.

The Periodic Table is printed on the back cover of this paper and the formulae for some common ions on the inside of the back cover.



JUN223430UE0101

Answer **all** questions.

1. Polymer gels are commonly used in disposable nappies.

A company that manufactures disposable nappies was investigating the effect of temperature on the mass of water the polymer gel in their nappies is able to absorb.

- (a) The results collected using water at 40 °C are given below. The initial mass of the polymer gel bead was 0.035 g.

Time (hours)	Mass of bead (g)	Mass of water absorbed by bead (g) (to 1 decimal place)
0	0.035	0.0
2	4.048	4.0
4	6.030	6.0
6	7.280	7.2
8	7.891	7.9
10	8.181	8.1
12	8.181	8.1

- (i) The percentage increase in the mass of the bead is calculated using the following equation.

$$\text{percentage increase} = \frac{\text{mass of water absorbed}}{\text{initial mass of bead}} \times 100$$

Calculate the percentage increase in the mass of the bead after 2 hours. Give your answer to the nearest whole number. [1]

Percentage increase = %


- (ii) What property of polymer gels does the figure calculated in part (i) demonstrate? [1]
-



- (b) (i) On the grid below, plot the results using water at 40 °C and draw a suitable line. Use the mass of water absorbed by the bead to 1 decimal place.

The results using water at 10 °C have already been plotted.

[3]



- (ii) Give **two** differences between the absorbing properties of the bead using water at 10 °C and at 40 °C.

[2]

Difference 1

.....

Difference 2


.....



2. The reactions of acids with metals, bases and carbonates are summarised in the following equations.

- acid + metal \rightarrow salt + hydrogen
- acid + base \rightarrow salt + water
- acid + carbonate \rightarrow salt + water + carbon dioxide

(a) The diagram shows some reactions of hydrochloric acid, HCl.



(i) Give the **names** of substances **A** and **B**. [2]

Substance **A**

Substance **B**

(ii) Give the **formulae** of substances **C** and **D**. [2]

Substance **C**


Substance **D**



(b) Complete the equation for the reaction between hydrochloric acid and sodium carbonate by

- writing the formula of sodium carbonate on the dotted line
- putting a number into the box to balance the equation

[2]





(c) Silver nitrate solution is used to identify the chloride ions present in hydrochloric acid.


(i) Give the observation made when silver nitrate solution is added to hydrochloric acid. [1]


.....


(ii) Put a tick (✓) in the box next to the correct ionic equation for the reaction between silver nitrate and hydrochloric acid. [1]














3. (a) The diagram shows the transfer of electrons that takes place during the formation of sodium oxide.



- (i) Name the type of bonding present in sodium oxide. [1]

.....

- (ii) State what must be done to sodium oxide so that it will conduct electricity.

Explain your answer.

[2]

.....
.....



- (b) (i) Draw a dot and cross diagram to show the bonding in a molecule of tetrafluoromethane, CF_4 . [2]

carbon (C) 2,4

fluorine (F) 2,7

- (ii) Tetrafluoromethane is a simple covalent substance and is a gas at room temperature.

Explain why it has a low boiling point. [2]

.....


.....

.....



- (c) Nanoparticles are extremely small particles that have different properties to the same material at bulk size.

Nano-scale titanium dioxide is commonly used in sunscreen and in self-cleaning windows.



Give the reason why nano-scale titanium dioxide is effective in each of these uses. [2]

Sunscreen

.....
.....

Self-cleaning windows

.....
.....



4. (a) When a mixture of iron(III) oxide and aluminium powder is heated, the following reaction takes place.



The reaction is commonly called the thermite reaction. The photographs show the reaction taking place and how it is used in the repair of railway lines.



- (i) Explain why iron is formed during the reaction. [2]

.....

.....

.....

- (ii) Complete and balance the equation for the reaction. [2]




- (iii) Calculate the percentage by mass of iron in iron(III) oxide, Fe_2O_3 . [2]

$$A_r(\text{Fe}) = 56 \quad A_r(\text{O}) = 16$$

Percentage = %



- (b) Clare and Frankie were investigating the reactivity of metals. They carried out a series of displacement reactions in a dropping tile. In each test they placed a small piece of metal into a solution of the nitrate of a different metal as shown.



- (i) It was not necessary to carry out all of the tests. Place crosses (×) on the diagram to show which tests did not need to be carried out.

Explain your choice.

[2]


.....

.....

.....



- (ii) The equation shows the reaction between iron and copper(II) nitrate solution.




Use the equation to calculate the maximum mass of copper that you would expect to be formed when 0.224 g of iron is added to excess copper(II) nitrate solution. [3]

$$A_r(\text{Fe}) = 56 \quad A_r(\text{Cu}) = 63.5$$

Maximum mass of copper = g




- (c) The photograph shows how the electrolysis of zinc chloride can be carried out in the laboratory.



- (i) Balance the equation that represents the reaction taking place at the anode.

Use the equation to explain the meaning of the term *oxidation*.

[2]



.....

.....

- (ii) The following method can be used to calculate the mass of zinc produced during the process.

- Record the mass of the cathode before placing it into the electrolyte
- Allow the process to run to completion
- Remove the cathode from the electrolyte and record its new mass
- Calculate the increase in mass

- I. It is often found that the increase in mass measured using this method is greater than expected. Suggest a reason for this.

[1]

.....

.....

- II. Suggest why this method cannot be used to measure the mass of chlorine produced during the process.

[1]

.....

.....

15




BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



5. (a) The bar chart shows the relative supply and demand for some fractions obtained from crude oil.



- (i) Use the chart to describe how the **difference** between supply and demand of the fractions changes as the chain length increases. [2]

.....

.....

.....

.....



(ii) The process of cracking is used to overcome the problem of insufficient supply of some fractions.

I. State what is meant by *cracking* and give the conditions needed for the process. [2]


.....

.....

.....

II. When hydrocarbon **X** is cracked, it forms hexene, ethene and butane.

Give the molecular formula of hydrocarbon **X** to complete the equation. [1]



(b) C_4H_{10} has two isomers.

(i) Give the meaning of the term *isomers*. [1]

.....

.....

.....

(ii) Draw the structures of both isomers of C_4H_{10} . [2]



6. It is well known that plastic bags are a major source of litter. These bags can exist for a very long time, ranging from tens of years in the natural environment up to a thousand years in landfill.

The use of plastic bags is now banned in some countries. These countries commonly use bags made from paper or cotton, which do not cause the same litter problem as plastic bags.

Paper or plastic?



Most people assume that a paper bag is better for the environment, however because paper bags are almost ten times heavier than plastic bags, they produce a greater mass of waste. On the other hand, because paper bags are biodegradable they do not lead to the same problem with litter as plastic bags.

When deciding whether paper is more suitable than plastic for making shopping bags, information from life cycle assessments (LCAs) of both materials must be taken into consideration.

Some of the findings from LCAs of paper and plastic bags are shown below.

Plastic bags are toxic to aquatic environments and their production is more likely to deplete natural resources

Plastic bags use 70% less energy in their production compared to paper bags

The processes used in paper production cause atmospheric acidification and eutrophication

The production of 1 million paper bags produces 51 tonnes of carbon dioxide, compared to 31 tonnes for plastic bags

693 kJ of energy is needed to recycle 1 kg of paper bags compared to 8 kJ for the same mass of plastic bags

4 546 litres of water are used to produce 1 000 paper bags compared to 177 litres for the same number of plastic bags

Paper bags are made from trees and plastic bags are made from ethene

Paper bags are 10 times heavier than plastic bags and therefore have more associated costs

On reading these findings it is important to consider all factors relating to sustainability when deciding on the most suitable material to make shopping bags. We should not just be thinking about what happens when we throw them away!



(a) Use the information in the passage to compare the mass of waste and volume of litter generated from the use of plastic and paper bags. [2]

.....
.....
.....
.....

(b) Use the information from the LCAs to tick (✓) whether each of the factors supports or opposes the use of paper bags in preference to plastic bags. [3]

	Supports use of paper bags	Opposes use of paper bags
Impact of waste on marine life	<input type="checkbox"/>	<input type="checkbox"/>
Water consumption in production	<input type="checkbox"/>	<input type="checkbox"/>
Energy used in production	<input type="checkbox"/>	<input type="checkbox"/>
Carbon footprint generated in production	<input type="checkbox"/>	<input type="checkbox"/>
Energy used in recycling	<input type="checkbox"/>	<input type="checkbox"/>
Cost of transporting waste	<input type="checkbox"/>	<input type="checkbox"/>

(c) The use of which type of bag is more likely to be linked to the erosion of limestone buildings? [1]

Give a reason for your answer.

Type of bag


Reason

.....

6



7. (a) The graph shows how the temperature changes when dilute hydrochloric acid is added gradually to sodium hydroxide solution.



The pH of the solution also changes as the hydrochloric acid is added to the sodium hydroxide.

Explain the temperature changes shown on the graph and relate these to the change in pH during the reaction. You do **not** need to include equations in your answer. [6 QER]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) The reaction between hydrochloric acid and sodium hydroxide produces sodium chloride and water.

Give the **ionic** equation to show how water is formed during this reaction. [1]

.....

END OF PAPER

7



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



FORMULAE FOR SOME COMMON IONS

POSITIVE IONS		NEGATIVE IONS	
Name	Formula	Name	Formula
aluminium	Al^{3+}	bromide	Br^-
ammonium	NH_4^+	carbonate	CO_3^{2-}
barium	Ba^{2+}	chloride	Cl^-
calcium	Ca^{2+}	fluoride	F^-
copper(II)	Cu^{2+}	hydroxide	OH^-
hydrogen	H^+	iodide	I^-
iron(II)	Fe^{2+}	nitrate	NO_3^-
iron(III)	Fe^{3+}	oxide	O^{2-}
lithium	Li^+	sulfate	SO_4^{2-}
magnesium	Mg^{2+}		
nickel	Ni^{2+}		
potassium	K^+		
silver	Ag^+		
sodium	Na^+		
zinc	Zn^{2+}		





THE PERIODIC TABLE

Group 1 2 3 4 5 6 7 0

		<table border="1"> <tr> <td style="text-align: center;">1</td> <td style="text-align: center;">H</td> <td style="text-align: center;">Hydrogen</td> <td style="text-align: center;">1</td> </tr> </table>		1	H	Hydrogen	1									<table border="1"> <tr> <td style="text-align: center;">4</td> <td style="text-align: center;">He</td> <td style="text-align: center;">Helium</td> <td style="text-align: center;">2</td> </tr> </table>		4	He	Helium	2																																						
1	H	Hydrogen	1																																																								
4	He	Helium	2																																																								
7	Li	Lithium	3	9	Be	Beryllium	4	11	Na	Sodium	11	12	Mg	Magnesium	12	19	F	Fluorine	9	20	Ne	Neon	10																																				
23	Na	Sodium	11	24	Mg	Magnesium	12	27	Al	Aluminium	13	28	Si	Silicon	14	31	P	Phosphorus	15	32	S	Sulfur	16	35.5	Cl	Chlorine	17	40	Ar	Argon	18																												
39	K	Potassium	19	40	Ca	Calcium	20	51	V	Vanadium	23	52	Cr	Chromium	24	56	Fe	Iron	26	59	Co	Cobalt	27	59	Ni	Nickel	28	63.5	Cu	Copper	29	65	Zn	Zinc	30	70	Ga	Gallium	31	73	Ge	Germanium	32	75	As	Arsenic	33	79	Se	Selenium	34	80	Br	Bromine	35	84	Kr	Krypton	36
86	Rb	Rubidium	37	88	Sr	Strontium	38	91	Zr	Zirconium	40	93	Nb	Niobium	41	96	Mo	Molybdenum	42	101	Ru	Ruthenium	44	103	Rh	Rhodium	45	106	Pd	Palladium	46	108	Ag	Silver	47	112	Cd	Cadmium	48	115	In	Indium	49	119	Sn	Tin	50	122	Sb	Antimony	51	127	I	Iodine	53	131	Xe	Xenon	54
133	Cs	Caesium	55	137	Ba	Barium	56	179	Hf	Hafnium	72	181	Ta	Tantalum	73	184	W	Tungsten	74	186	Re	Rhenium	75	192	Ir	Iridium	77	195	Pt	Platinum	78	197	Au	Gold	79	201	Hg	Mercury	80	204	Tl	Thallium	81	207	Pb	Lead	82	210	Po	Polonium	84	210	At	Astatine	85	222	Rn	Radon	86
223	Fr	Francium	87	226	Ra	Radium	88	227	Ac	Actinium	89	227	La	Lanthanum	57	227	Ac	Actinium	89																																								

Key

